


x a Niche company

Hillandale

Unit 1, 53 Southgate Avenue Cannon Hill

Brisbane, QLD 4170 Phone: +61 407 332 063 E-mail: info@ausecology.com Web: www.ausecology.com



Prepared for: Report issued: Senex August 2025

Senex

ATLAS Stage 3 — EPBC Pre-clearance Report - Hillandale August 2025



This document has been prepared and is certified by: AUSECOLOGY PTY LTD

ABN 15 155 304 751

PO Box 594, Morningside, QLD 4170

w www.ausecology.com e info@ausecology.com

Document status

| Revision | Reason for issue | Author | Reviewed | Issued to | Date |
|----------|------------------------------|------------------------|----------|--------------------|------------|
| А | Draft report | B. Osborne M. Neill | - | - | 13/03/2024 |
| 0 | Final Report | B. Osborne M. Neill | L. Eade | Senex | 14/03/2025 |
| 1 | Final Report | B. Osborne M. Neill | L. Eade | Senex | 18/03/2025 |
| 2 | Updated mapping | B. Osborne M. Neill | L. Grigg | Senex | 21/03/2025 |
| 3 | Issue to client after review | B. Osborne M. Neill | L. Grigg | P. Wilkison -Senex | 09/05/2025 |
| 4 | Issue to client after review | B. Osborne M. Neill | L. Grigg | P. Wilkison -Senex | 14/08/2025 |



Table of contents

| Glo | ssary of | f Terms | iv |
|------|-----------|---|---------------------------------------|
| 1 | Intro | duction | 1 |
| | 1.1 | Project background | 1 |
| | 1.2 | Scope | |
| | 1.3 | EPBC conditions | |
| 2 | Meth | nodology | 2 |
| | 2.1 | Regional ecosystem assessment and threatened ecological communities | 2 |
| | 2.2 | Targeted threatened flora surveys | 2 |
| | 2.3 | Fauna habitat assessment | 3 |
| | 2.4 | Threatened fauna surveys | |
| | 2.5 | Koala dispersal habitat | |
| | 2.6 | Squatter pigeon dispersal habitat | |
| 3 | Resul | lts | 6 |
| | 3.1 | Regional ecosystems and threatened ecological communities | 6 |
| | 3.2 | Targeted threatened flora surveys | 6 |
| | 3.3 | Opportunistic fauna survey and habitat assessment | 6 |
| | 3.4 | Weeds | |
| | 3.5 | Erosion | 7 |
| | 3.6 | Squatter pigeon habitat | 7 |
| | 3.7 | Ground-truthed koala dispersal trees | 7 |
| Disc | cussion_ | | 48 |
| App | endix A | A – Fauna records | 49 |
| App | endix E | 3 – Habitat features | 51 |
| App | endix C | C – Weed species list | 54 |
| App | endix [| D – Erosion points and photos | 56 |
| | | | |
| Ta | bles | | |
| Tab | le 3-1 | Dispersal habitat trees | 8 |
| Fig | ures | | |
| Figu | ıre 2-1 (| Overview Map | 5 |
| | ıre 3-1 | Representative image of cleared grazing pastures | |
| Figu | ıre 3-2 | Representative image of patchy regrowth vegetation | |
| _ | ıre 3-3 | Scattered course woody debris between Atlas-274 and Atlas-253 | |
| _ | ıre 3-4 | Numerous scattered stags south of Atlas-258 | |
| _ | ire 3-5 | Ground-truthed Koala Tree Extents Mapping | |
| ۰, | | | · · · · · · · · · · · · · · · · · · · |



Glossary of Terms

| Acronym | Description |
|---------------|--|
| ATP | Authority to prospect |
| DBH | Diameter at breast height |
| EA | Environmental authority |
| ECPPFD | Environmental Constraints Protocol for Planning and Field Development |
| EPBC Act | Environment Protection and Biodiversity Conservation Act 1999 (Commonwealth) |
| EP Regulation | Environmental Protection Regulation 2008 |
| ESA | Environmentally Sensitive Area |
| GTRE | Ground-truthed regional ecosystem |
| ha | Hectare |
| km | Kilometres |
| m | Metres |
| MNES | Matter of National Environmental Significance |
| MSES | Matter of State Environmental Significance |
| PL | Petroleum lease |
| sp. | Species (singular) |
| spp. | Species (plural) |
| Sqm | Square metres |
| SQP | Suitably qualified person |
| TEC | Threatened Ecological Community |



1 Introduction

The pre-clearance survey methodology applied within this package of works is deemed appropriate to confirm the on-the-ground biodiversity values present.

1.1 Project background

The Atlas Stage 3 Gas Project (EPBC Act referral 2022/09410) involves developing, operating, decommissioning and rehabilitating up to 151 coal seam gas wells; gas and water gathering systems for the producing wells; access tracks; brine and produced water storages; borrow pits; and ancillary supporting facilities on Petroleum Lease (PL) 1127, PL1037, PL445 and PL209 in the central part of the Surat Basin, Queensland (Senex, 2024).

This report provides the results of pre-clearance survey on a proposed infrastructure layout including wells, gathering, camp, laydowns and extra workspaces (known hereafter as 'The Footprint') and a 30 m buffer within the Hillandale property:

- Lot 50 on Plan FT167
- Lot 51 on Plan FT429

1.2 Scope

Ausecology Pty Ltd (Ausecology) was engaged by Senex Energy Pty Ltd (Senex) to undertake pre-clearance ecological surveys as part of the approval conditions for the Atlas Stage 3 Gas Project and in accordance with the Atlas Stage 3 Environmental Constraints Protocol for Planning and Field Development (ECPPFD) document (Senex, 2024). The ECPPFD provides a framework for identifying, assessing and managing potential impacts to Matters of National Environmental Significance (MNES) and Matter of State Environmental Significance (MSES) associated with development of the Atlas Stage 3 Gas Project. Data collected during the pre-clearance surveys will be used by Senex to ensure:

- infrastructure siting complies with relevant environmental approval conditions and does not exceed the maximum disturbance limits
- infrastructure siting adheres to the constraints mapping; and
- no functional change to Koala dispersal habitat, the approval holder must not remove more than a total of 4 ha of trees, measured in canopy cover within mapped koala dispersal habitat.

Results from the pre-clearance survey findings (this report) will be published on the website, including:

- the location and extent of trees to be cleared, including maps; and
- a discussion of how removal of trees will not change the ability of koalas to disperse across the landscape.

This survey also involved targeted threatened flora surveys, active threatened fauna surveys (where suitable habitat was identified) and fauna habitat identification (where encountered) as per the ECPPFD.

1.3 EPBC conditions

This report will validate compliance with the following EPBC approval conditions:

- 1. The approval holder must not:
 - a) clear any Koala foraging and breeding habitat.
 - b) clear more than 2.1 hectares (ha) of Squatter Pigeon dispersal habitat.



- 4. In accordance with the Constraints Protocol, the approval holder must:
 - b) adhere to the constraints mapping.
 - 5. To ensure no functional change to Koala dispersal habitat, the approval holder must not remove more than a total of 4 ha of trees, measured in canopy cover within mapped Koala dispersal habitat.

PRE-CLEARANCE SURVEYS

- 6. Prior to commencing clearing, the approval holder must:
 - a) undertake at least one pre-clearance survey of the proposed area of clearing, and
 - b) publish on the website the pre-clearance survey findings, including:
 - i) the location and extent of trees to be cleared, including maps, and
 - *ii)* a discussion of how removal of trees will not change the ability of Koalas to disperse across the landscape.

2 Methodology

An Ausecology senior ecologist (suitably qualified) and ecologist conducted field surveys on foot on the 3rd to the 6th of March 2025 to determine impacts from the planned gas infrastructure including wells, gathering, laydowns extra workspaces and a camp on the Hillandale property. The area surveyed and mapped in this report is shown in Figure 2-1.

2.1 Regional ecosystem assessment and threatened ecological communities

2.1.1 Desktop assessment

Desktop assessment and constraints assessments have previously been undertaken on the property including by Ausecology in November 2024). These surveys included ground-truthed regional ecosystem (GTRE) mapping and threatened ecological community surveys. This report has been reviewed as part of this desktop assessment prior to the pre-clearance surveys and were deemed to have been conducted to an acceptable level of detail. Given the level of detail in prior reports, no further detailed desktop analysis has been conducted.

2.1.2 Regional ecosystem assessment

During pre-clearance surveys, quaternary site assessments to verify regional ecosystems were undertaken where necessary (i.e., where vegetation and ecological communities have been determined to vary from the mapped GTRE at the time of the pre-clearance surveys). These assessments were conducted in accordance with the ECPPFD.

2.1.3 Threatened ecological communities

Where necessary Threatened Ecological Community (TEC) assessments were undertaken to confirm the presence and condition of TECs identified as known or potential in The Footprint if these were determined to vary from the mapped TEC areas identified in the constraints mapping.

2.2 Targeted threatened flora surveys

These surveys were conducted by a suitably qualified person (SQP). Targeted flora surveys of all known, likely or potential threatened flora species were conducted within The Footprint and 30 m buffer, where mapped constraint areas were present and/or suitable habitat was identified by the SQP, in accordance with the ECPPFD.

These surveys were conducted using the random meander methodology and if a species was encountered, a population survey was undertaken to determine the extent and density of the population. Threatened flora

Senex

ATLAS Stage 3 – EPBC Pre-clearance Report - Hillandale August 2025



species and the locations of all individuals were recorded and specimens collected of any unknown individual requiring confirmation by the Queensland Herbarium.

Potentially occurring threatened species in the area as per the ECPPFD include Belson's panic (*Homopholis belsonii*), red soil woolly wrinklewort (*Rutidosis lanata*) and winged nightshade (*Solanum stenopterum*).

2.3 Fauna habitat assessment

Senex has committed to not clearing any areas confirmed as habitat for threatened species (ECPPFD), with the exception of koala (*Phascolarctos cinereus*) dispersal habitat and short-beaked echidna (*Tachyglossus aculeatus*) habitat. Fauna habitat baseline assessments have been conducted to an adequate level of detail to enable known, likely and potentially present species to be identified and comprehensive Project impact assessment and constraints mapping has been completed (ECPPFD).

The pre-clearance surveys will reassess the habitat present (as mapped in the constraints mapping (ECPPFD)) or otherwise identified by the SQP during the pre-clearance surveys) in order to refine mapped habitat areas. They will also identify and record micro-habitat features and breeding sites to facilitate avoidance and minimisation of impacts to potentially utilised micro-habitat features and breeding sites. Recorded micro-habitat features, where present include:

- Hollow-bearing trees;
- Dead standing trees;
- Hollow logs;
- Termite mounds;
- Woody debris;
- Surface rocks;
- Gilgais;
- Soil cracks / cracking clay;
- Rocky outcrops, crevices, overhangs and caves;
- Mistletoes;
- Nests;
- Animal burrows;
- Watercourses, wetlands and dams (including proximity); and
- Any other significant habitat features, or values present, such as dense leaf litter, heavily decorticating bark, dense grass/shrub shelter, seeding grass cover, fruiting plants, nectar and pollen producing plants and koala food trees.

2.4 Threatened fauna surveys

As areas confirmed as habitat for threatened species have been effectively avoided by The Footprint (with the exception of koala dispersal habitat and echidna habitat), the area required to be surveyed was minimal.

Active fauna surveys of all known, likely or potential threatened fauna species are to be undertaken where suitable potential habitat is mapped or found to be present within The Footprint (refer to the constraints mapping and the habitat features listed in Appendix A of the ECPPFD). Active fauna searches as per Table 1, Appendix A of the ECPPFD include scanning trees, the ground and habitat features; overturning rocks, logs and

Senex

ATLAS Stage 3 – EPBC Pre-clearance Report - Hillandale August 2025



other woody debris; searching under peeling bark; raking leaf litter and soil at the base of trees; and flushing birds from dense shrubs and groundcover.

Invasive active searches were not undertaken in the 30 m buffer of The Footprint within mapped constraint areas, given no impact is expected and active searches outside of The Footprint would be detrimental to the fauna species habitat. Instead of invasive searches in the 30 m buffer, surveys included incidental observations and scat and sign searches.

2.5 Koala dispersal habitat


An initial desktop survey was undertaken to analyse all previous ecology data collected in the field, analysis of the ECPPFD and associated constraints mapping, and to identify areas of mapped koala dispersal habitat that would require further ground verification.

Areas identified were highlighted on GIS mapping for further identification and field verification. Additional points were collected in the field where applicable. All tree ID numbers have been provided in individual maps of the area surveyed and further details provided in the results.

Canopy cover was measured by walking the dripline of trees located in koala dispersal areas using a sub-10 cm accuracy handheld Trimble GPS unit. The diameter at breast height (DBH) of each of the abovementioned trees was measured and height data was collected. Where denser regrowth was present and canopy cover of individual trees could not be distinguished, the canopy cover of the clump of trees was walked at the outer dripline and average height assessed. Thorny tree species which are not able to be climbed by koala (i.e. desert lime (*Citrus glauca*) were noted in the field but excluded from the final koala tree canopy cover calculations.

2.6 Squatter pigeon dispersal habitat

An initial desktop survey was undertaken to analyse all previous ecology data collected in the field, analysis of the ECPPFD and associated constraints mapping, and to identify areas of mapped squatter pigeon dispersal habitat that would require further ground verification.





3 Results

The Hillandale property has been historically cleared for cattle grazing and is dominated by cleared pasture lands of introduced grasses and scattered paddock trees such as poplar box (*Eucalyptus populnea*), brigalow (*Acacia harpophylla*), belah (*Casuarina cristata*) and wilga (*Geijera parviflora*) (Figure 3-2). Small patches of remnant and regrowth vegetation were also found throughout the property (Figure 3-2).





Figure 3-1 Representative image of cleared Figure 3-2 Representative image of patchy grazing pastures regrowth vegetation

3.1 Regional ecosystems and threatened ecological communities

Where mapped constraints areas occurred within the disturbance footprint or 30 m buffer these were assessed in the field to ensure they aligned with the mapping. No mapped constraints in the survey area were found to require changes. The survey also confirmed that the proposed disturbance areas do not contain any listed TEC or any TEC within the 30 m buffer area.

3.2 Targeted threatened flora surveys

No Belson's panic (*Homopholis belsonii*), red soil woolly wrinklewort (*Rutidosis lanata*) and winged nightshade (*Solanum stenopterum*) were identified within The Footprint. It is unlikely for these species to be present in The Footprint, due to the grazing pressure and presence of non-native grasses, mainly buffel grass (*Cenchrus ciliaris*).

3.3 Opportunistic fauna survey and habitat assessment

No threatened fauna species were observed in suitable potential habitat within 30 m of The Footprint, a total of 31 incidental fauna species were found, with the full list shown in Appendix A. Habitat searches found 20 microhabitat features other than koala dispersal trees including course woody debris, dead standing trees (stags), decorticating bark and bird nests within The Footprint and 30 m buffer, with a full list shown in Appendix B. All nests identified in The Footprint did not have any nesting birds at the time of the survey. Invasive searches were not undertaken on habitat features in the proposed footprint, as fauna would be disturbed too soon before clearing, and habitat may be unnecessarily destroyed.

Within The Footprint two polygons were mapped as habitat due to the large number of features present. One area mapped between Atlas-274 and Atlas-253 had frequent course woody debris and logs from past clearing (Figure 3-3). The second mapped polygon south of Atlas-258 has numerous stags with some containing hollows, cracks or decorticating bark as well as some course woody debris (Figure 3-4).







Figure 3-3 Scattered course woody debris between Figure 3-4
Atlas-274 and Atlas-253

Figure 3-4 Numerous scattered stags south of Atlas-258

3.4 Weeds

Restricted invasive species under the *Biosecurity Act 2014* present within The Footprint included scattered *Opuntia stricta* and *Opuntia tomentosa*. An additional 18 weed species were also noted, see Appendix C for a list of weeds identified in The Footprint.

3.5 Erosion

Multiple erosion points were recorded around the property, particularly along drainage lines. Several erosion issues in proximity to The Footprint are shown in Appendix D.

3.6 Squatter pigeon habitat

A total of 331.6 ha of Squatter pigeon habitat was mapped across the Hillendale site. All breeding and foraging habitat for the species has been avoided and a total of 0.18 ha of dispersal habitat for the species to be impacted as a result of the Project (Figure 3-26). This is within the 2.1 ha of allowable clearing limits for squatter pigeon dispersal habitat.

3.7 Ground-truthed koala dispersal trees

Table 3-1 provides the results including canopy cover, height and DBH of the trees assessed in the field in koala dispersal areas. All trees were assessed and, where determined they could be avoided (through underground boring or realignment), have been marked up as "retain" and will be avoided during construction. Trees unable to be avoided have had their canopy cover assessed and calculated under disturbance limits. Mapping of each location (Desktop ID) are shown in Figure 3-5 to Figure 3-25.



Table 3-1 Dispersal habitat trees

| Location | Area | l habitat trees | DBH | Height | | |
|--------------|-------|------------------------|------|--------|--------|-------|
| (Desktop ID) | (sqm) | Species | (cm) | (m) | Action | Photo |
| HLLO1 | 48.48 | Geijera parviflora | 39 | 8.2 | Retain | |
| HLL02 | 14.20 | Casuarina cristata | 10 | 5.3 | Remove | |
| HLL03 | 13.59 | Owenia acidula | 23 | 5.8 | Retain | |
| HLLO4 | 35.02 | Callitris glaucophylla | 40 | 12.5 | Retain | |
| HLL05 | 78.40 | Callitris glaucophylla | 31 | 11.5 | Retain | |
| HLLO6 | 7.73 | Callitris glaucophylla | 14 | 6.3 | Retain | |
| HLL07 | 5.19 | Callitris glaucophylla | 10 | 6.2 | Retain | |
| HLL08 | 8.80 | Callitris glaucophylla | 15 | 6.5 | Retain | |



| Location (Desktop ID) | Area (sqm) | Species | DBH (cm) | Height (m) | Action | Photo |
|--------------------------|---------------|----------------------------|-------------|---------------|--------|-------|
| HLL09 | 5.22 | Callitris glaucophylla | 14 | 4.4 | Retain | |
| HLL10 | 11.20 | Acacia pendula | 20 | 6.5 | Retain | |
| HLL11 | 9.28 | Callitris glaucophylla | 16 | 6 | Retain | |
| HLL12 | 21.30 | Casuarina cristata | 22 | 10.2 | Retain | |
| HLL13 | 38.46 | Eucalyptus melanophloia | 32 | 9.9 | Remove | |
| HLL14 | 56.40 | Eucalyptus melanophloia | 23 | 10.1 | Kemove | |
| HLL15 | 22.82 | Acacia excelsa | 10 | 4.5 | Remove | |
| HLL16 | 22.02 | Eucalyptus melanophloia | 30 | 10.1 | Nemove | |
| HLL19 | 42.51 | Geijera parviflora | 37 | 5.6 | Retain | |



| Location (Desktop ID) | Area (sqm) | Species | DBH (cm) | Height (m) | Action | Photo |
|--------------------------|---------------|----------------------------|-------------|---------------|--------|-------|
| HLL20 | 30.63 | Geijera parviflora | 28 | 6.6 | Retain | |
| HLL21 | 76.29 | Geijera parviflora | 47 | 11.7 | Retain | |
| HLL22 | 3.80 | Eucalyptus melanophloia | 10 | 4.7 | Retain | |
| HLL23 | 2.92 | Eucalyptus melanophloia | 10 | 4.7 | Retain | |
| HLL24 | 58.55 | Geijera parviflora | 41 | 8 | Retain | |
| HLL25 | 76.09 | Acacia harpophylla | 16 | 9.4 | Retain | |
| HLL26 | 38.60 | Acacia harpophylla | 18 | 7.4 | Retain | |
| HLL28 | 6.99 | Acacia harpophylla | 12 | 6.6 | Retain | |



| Location (Desktop ID) | Area (sqm) | Species | DBH (cm) | Height (m) | Action | Photo |
|--------------------------|---------------|---------------------|-------------|---------------|--------|-------|
| HLL29 | 20.78 | Acacia harpophylla | 18 | 6.6 | Retain | |
| HLL30 | 8.06 | Acacia harpophylla | 18 | 6.8 | Retain | |
| HLL33 | 15.07 | Atalaya hemiglauca | 21 | 7.2 | Retain | |
| HLL34 | 23.22 | Geijera parviflora | 20 | 5.3 | Remove | |
| HLL35 | 4.69 | Eucalyptus populnea | 25 | 8 | Retain | |
| HLL36 | 10.38 | Acacia harpophylla | 15 | 6.6 | Retain | |
| HLL37 | 76.35 | Acacia salicina | 30 | 7 | Remove | |
| HLL38 | 28.48 | Acacia salicina | 15 | 6 | Retain | |



| Location (Desktop ID) | Area (sqm) | Species | DBH (cm) | Height (m) | Action | Photo |
|--------------------------|---------------|------------------------|-------------|---------------|--------|-------|
| HLL39 | 30.30 | Acacia salicina | 21 | 6.8 | Retain | |
| HLL40 | 5.20 | Eremophila mitchellii | 18 | 6.6 | Remove | |
| HLL41 | 11.45 | Eucalyptus orgadophila | 21 | 9.8 | Retain | |
| HLL42 | 64.64 | Brachychiton rupestris | 106 | 7.1 | Remove | |
| HLL43 | 9.22 | Eucalyptus populnea | 17 | 6.4 | Retain | |
| HLL44 | 18.19 | Eremophila mitchellii | 23 | 7.3 | Retain | |
| HLL45 | 11.85 | Geijera parviflora | 9 | 5 | Retain | |



| Location (Desktop ID) | Area (sqm) | Species | DBH (cm) | Height (m) | Action | Photo |
|--------------------------|---------------|---------------------|-------------|---------------|--------|-------|
| HLL46 | 5.60 | Eucalyptus populnea | 10 | 5.5 | Retain | |

| HLL47 | 14.43 | Acacia harpophylla | 16 | 5.8 | Retain | |
|-------|--------|-----------------------|----|------|--------|--|
| HLL48 | 20.04 | Acacia harpophylla | 28 | 9.3 | Retain | |
| HLL49 | 19.10 | Acacia harpophylla | 21 | 5.7 | Retain | |
| HLL50 | 110.19 | Owenia acidula | 42 | 9.1 | Retain | |
| HLL51 | 33.04 | Eremophila mitchellii | 16 | 6.6 | Retain | |
| HLL52 | 10.48 | Geijera parviflora | 14 | 4.3 | Remove | |
| HLL53 | 48.15 | Owenia acidula | 40 | 9 | Remove | |
| HLL54 | 234.65 | Owenia acidula | 65 | 13.8 | Retain | |



| Location (Desktop ID) | Area (sqm) | Species | DBH (cm) | Height (m) | Action | Photo |
|--------------------------|---------------|--------------------|-------------|---------------|--------|-------|
| HLL56 | 21.14 | Acacia salicina | 25 | 7 | Retain | |
| HLL57 | 22.48 | Geijera parviflora | 15 | 6.3 | Retain | |
| HLL59 | 6.37 | Acacia harpophylla | 11 | 6 | Retain | |
| HLL61 | 22.99 | Acacia salicina | 11 | 7.2 | Retain | |
| HLL62 | 107.45 | Acacia salicina | 38 | 11 | Retain | |
| HLL63 | 35.77 | Acacia salicina | 35 | 12.5 | Retain | |
| HLL64 | 30.01 | Acacia harpophylla | 36 | 7.1 | Retain | |



| Location (Desktop ID) | Area (sqm) | Species | DBH (cm) | Height (m) | Action | Photo |
|--------------------------|---------------|----------------------|-------------|---------------|--------|-------|
| HLL65 | 9.62 | Acacia harpophylla | 44 | 14.6 | Retain | |
| HLL66 | | Acacia harpophylla | 29 | 13.4 | | |
| HLL67 | 4.44 | Acacia harpophylla | 24 | 10.8 | Retain | |
| HLL68 | 12.15 | Acacia harpophylla | 32 | 9.8 | Retain | |
| HLL69 | 8.86 | Santalum lanceolatum | 19 | 7.3 | Retain | |
| HLL70 | 18.83 | Acacia salicina | 31 | 9.2 | Retain | |
| HLL71 | 57.47 | Acacia salicina | 20 | 6.8 | Retain | |
| HLL72 | 41.70 | Acacia salicina | 27 | 8.5 | Retain | |
| HLL73 | 6.16 | Acacia salicina | 27 | 9.8 | Retain | |
| HLL74 | 3.04 | Acacia salicina | 17 | 4.5 | Retain | |
| HLL75 | 8.2 | Acacia salicina | 15 | 4 | Retain | N/A |



| Location (Desktop ID) | Area (sqm) | Species | DBH (cm) | Height (m) | Action | Photo |
|--------------------------|---------------|----------------------|-------------|---------------|--------|-------|
| HLL76 | 10.79 | Acacia salicina | 11 | 4.7 | Retain | |
| HLL77 | 23.56 | Acacia salicina | 27 | 27 8.5 Retain | | |
| HLL78 | 19.40 | Acacia salicina | 30 | 5.3 | Retain | |
| HLL79 | 16.48 | Acacia salicina | 7.4 | 35 | Retain | |
| HLL80 | 9.80 | Acacia salicina | 17 | 9.5 | Retain | |
| HLL82 | 11.38 | Geijera parviflora | 10 | 4.5 | Retain | |
| HLL83 | 14.67 | Atalaya hemiglauca | 23 | 23 7 Retain | | |
| HLL84 | 27.30 | Geijera parviflora | 16 | 5.9 | Retain | |
| HLL85 | 9.12 | Santalum lanceolatum | 14 | 4.9 | Retain | |



| Location (Desktop ID) | Area (sqm) | Species | DBH (cm) | Height (m) | Action | Photo |
|--------------------------|---------------|-----------------------|-------------|---------------|--------|------------|
| HLL86 | 109.61 | Owenia acidula | 46 | 12.1 | Retain | |
| HLL87 | 34.67 | Geijera parviflora | viflora 48 | | Retain | |
| HLL88 | 9.54 | Acacia salicina | 12 | 5 | Retain | |
| HLL89 | 34.27 | Acacia salicina | 15 | 6.2 | Retain | |
| HLL90 | 11.74 | Acacia salicina | 11 | 4.5 | Retain | |
| HLL91 | 12.22 | Acacia salicina | 14 | 6.5 | Retain | |
| HLL92 | 7.90 | Acacia salicina | 20 | 8 | Retain | All Market |
| HLL93 | 3.45 | Eremophila mitchellii | 22 | 6.2 | Retain | |



| Location (Desktop ID) | Area (sqm) | Species | DBH (cm) | Height (m) | Action | Photo |
|--------------------------|---------------|------------------------|-------------|---------------|--------|-------|
| HLL94 | 21.03 | Acacia harpophylla | 14 | 4.9 | Remove | |
| HLL95 | 71.52 | Acacia harpophylla | 24 | 24 9.6 Retain | | |
| HLL96 | 39.12 | Acacia harpophylla | 24 | 24 5.8 Retain | | |
| HLL97 | 3.53 | Acacia harpophylla | 12 | 4.8 | Remove | |
| HLL98 | 36.61 | Atalaya hemiglauca | 13 | 13 5.3 Remove | | |
| HLL99 | 11.10 | Acacia excelsa | 15 | 6.1 | Remove | |
| HLL100 | 8.07 | Eucalyptus orgadophila | 10 | 5.5 | Remove | |



| Location (Desktop ID) | Area (sqm) | Species | DBH Height (cm) (m) Action | | Action | Photo |
|--------------------------|---------------|------------------------|----------------------------|----------------|--------|-------|
| HLL101 | 79.03 | Acacia salicina | 30 | 30 13.3 Retain | | |
| HLL102 | 16.12 | Atalaya hemiglauca | 27 | 9.1 | Remove | |
| HLL103 | 8.29 | Atalaya hemiglauca | 25 | 9 | Remove | |
| HLL104 | 47.43 | Brachychiton rupestris | 190 | 9.5 | Retain | |
| HLL105 | 48.39 | Brachychiton rupestris | 180 | 11.4 | Retain | |
| HLL106 | 19.19 | Brachychiton rupestris | 150 | 150 8.7 Retain | | |
| HLL107 | 31.99 | Acacia harpophylla | 15 | 6.5 | Retain | |
| HLL108 | 23.03 | Acacia harpophylla | 17 | 8.7 | Remove | |



| Location (Desktop ID) | Area (sqm) | Species | DBH (cm) | Height (m) | Action | Photo |
|--------------------------|---------------|------------------------|---------------|-----------------|--------|-------|
| HLL109 | 8.79 | Acacia harpophylla | 12 | 5.3 | Remove | |
| HLL110 | 6.02 | Acacia harpophylla | 14 5.2 Remove | | Remove | |
| HLL112 | 33.60 | Brachychiton rupestris | 100 | 100 10.9 Retain | | |
| HLL113 | 16.13 | Brachychiton rupestris | 70 | 6 | Retain | |
| HLL114 | 38.54 | Acacia harpophylla | 13 | 13 5.2 Remove | | |
| HLL115 | 11.14 | Acacia harpophylla | 15 | 7.1 | Remove | |
| HLL116 | 20.04 | Acacia harpophylla | 12 | 5.2 | Retain | N/A |
| HLL117 | 16.91 | Notelaea macrocarpa | 14 | 4.2 | Retain | |



| Location (Desktop ID) | Area (sqm) | Species | DBH (cm) | Height (m) | Action | Photo |
|--------------------------|---------------|------------------------|----------------|----------------|--------|-------|
| HLL118 | 5.61 | Acacia harpophylla | 10 | 10 4.7 | | |
| HLL119 | 33.04 | Acacia harpophylla | 20 | 8 | Retain | |
| HLL121 | 28.57 | Brachychiton rupestris | 170 9.1 Retain | | Retain | |
| HLL122 | 37.35 | Acacia salicina | 10 | 10 5.3 Retain | | |
| HLL123 | 35.69 | Acacia salicina | 10 | 7.3 | Retain | |
| HLL124 | 108.21 | Owenia acidula | 63 | 63 12.9 Remove | | |
| HLL126 | 47.81 | Acacia harpophylla | 30 | 8 | Retain | |
| HLL127 | 38.78 | Acacia harpophylla | 30 | 8 | Remove | |



| Location (Desktop ID) | Area (sqm) | Species | DBH (cm) | Height (m) | Action | Photo |
|--------------------------|---------------|---------------------|-------------|---------------|--------|-------|
| HLL128 | 21.85 | Eucalyptus populnea | 32 9 Remove | | Remove | |
| HLL129 | 10.75 | Acacia harpophylla | 15 | 5.5 | Remove | |
| HLL131 | 44.08 | Acacia harpophylla | 14 | 6.1 | Retain | |
| HLL132 | 10.11 | Casuarina cristata | 13 | 6.1 | Retain | |
| HLL133 | 14.82 | Casuarina cristata | 12 | 7.6 | Retain | |
| HLL134 | 19.19 | Casuarina cristata | 12 | 6.8 | Retain | |
| HLL135 | 10.01 | Casuarina cristata | 10 | 5.7 | Retain | |



| Location (Desktop ID) | Area (sqm) | Species | DBH (cm) | Height (m) | Action | Photo |
|--------------------------|---------------|--------------------|-------------|---------------|--------|----------|
| HLL136 | 6.58 | Casuarina cristata | 10 | 6 | Retain | |
| HLL137 | 7.89 | Acacia harpophylla | 12 | 12 5.8 Retain | | |
| HLL140 | 15.43 | Geijera parviflora | 18 | 5.8 | Retain | |
| HLL141 | 43.95 | Geijera parviflora | 68 | 8.3 | Retain | |
| HLL142 | 3.50 | Acacia salicina | 10 | 10 4.5 Reta | | Live con |
| HLL143 | 10.73 | Geijera parviflora | 12 | 7 | Retain | |
| HLL144 | 35.21 | Geijera parviflora | 17 | 5 | Retain | |




| Location (Desktop ID) | Area (sqm) | Species | DBH (cm) | Height (m) | Action | Photo |
|--------------------------|---------------|-----------------------|------------------------|-------------|--------|-------|
| HLL145 | 7.10 | Geijera parviflora | 10 | 4.7 | Retain | |
| HLL146 | 9.70 | Geijera parviflora | 20 | 6 | Retain | N/A |
| HLL147 | 5.31 | Geijera parviflora | 15 | 6 | Retain | N/A |
| HLL149 | 2.99 | Eucalyptus populnea | ptus populnea 11 8 Rem | | Remove | |
| HLL150 | 17.08 | Geijera parviflora | 10 4.8 Retain | | Retain | |
| HLL151 | 40.34 | Eremophila mitchellii | 15 | 15 6.1 Reta | | |
| HLL152 | 12.98 | Eucalyptus populnea | 15 | 15 6.6 | | |
| HLL153 | 36.96 | Geijera parviflora | 34 | 7.9 | Remove | |





| Location (Desktop ID) | Area (sqm) | Species | | DBH (cm) | Height (m) | Action | Photo |
|--------------------------|---------------|-------------------------------|-------------------|-------------|---------------|--------|--|
| HLL154 | 2.42 | Casuarina cristata | | 10 | 7.9 | Retain | |
| HD46 | 26.12 | Eucalyptus po | pulnea | 20 | 5 | Retain | N/A |
| HLL155 | 4.94 | | | | | Retain | |
| HLL156 | 1.51 | | | | | Retain | |
| HLL157 | 1.47 | | | | Retain | | |
| HLL158 | 2.69 | | | | Retain | | |
| HLL159 | 1.52 | | | | | Retain | |
| HLL160 | 1.50 | | | | | Retain | |
| HLL161 | 1.71 | | | | | Retain | |
| HLL162 | 1.27 | | | | | Retain | and the second s |
| HLL163- HLL164 | 1.46 | Patch of regr Acacia salid | Patch of regrowth | | | Retain | |
| HLL165 | 2.71 | | | Retain | | | |
| HLL166 | 1.23 | | Retain | | | | |
| HLL167- HLL168 | 15.10 | 10 | | Retain | | | |
| HLL169 | 7.24 | | | | | Retain | |
| HLL170 | 4.80 | | | | | Retain | |
| HLL171 | 5.83 | | | | | Retain | |
| HLL172 | 15.88 | | | | | Retain | |
| HLL173 | 13.16 | | | | | Retain | |
| HLL174 | 21.63 | | | | | Retain | |
| HLL175 | 16.22 | | | | | Retain | |
| HLL176 | 29.05 | | | | | Retain | |
| HLL177 | 12.87 | | | | | Retain | |
| HLL178- HLL181 | 39.32 | Patch of regr | | Avg: | | Remove | A Company of the Comp |
| HLL182- HLL184 | 28.68 | scattered Ge parviflor | eijera | 15 | Avg: 5 | Remove | 4,14 |
| HLL185- HLL187 | 13.46 | | | | | Remove | |
| HLL188 | 31.91 | 1 | | | | Remove | |
| HLL189 | 9.66 | | | | | Retain | |
| HLL190 | 85.58 | | | | | Retain | |
| Remove area: | | Total (sqm) | 828.69 | | | | |
| | | Total (ha) | 0.082 | | | | |
| Retain area: | | Total (sqm) | 3181.74 | ! | | | |
| | | Total (ha) | 0.32 | | | | |
























































ATLAS Stage 3 – EPBC Pre-clearance Report - Hillandale August 2025



Discussion

The surveys on the Hillandale property found that the projects' impacts to koala and squatter pigeon dispersal habitat would be minimal with a total potentially to be cleared of 0.082 ha and 0.18 ha respectively. The majority of the impact footprint is located within predominantly cleared agricultural areas and has actively avoided major remaining fragments of habitat and potential koala dispersal trees following the constraint mapping. Trees retained within close proximity to those that are unable to be avoided total 0.32 ha, and the removal of 0.082 ha is unlikely to have a significant effect on the ecological function of dispersal habitat on the property for koalas.

No remnant or HVR regulated vegetation, TEC and potential threatened fauna habitat are present within The Footprint. Threatened flora searches found no threatened species within or in proximity to The Footprint. Numerous weed species were observed on the property, notably listed restricted and WoNS; common prickly pear (*Opuntia stricta*) and velvety tree pear (*Opuntia tomentosa*).

Numerous other least concern species were observed in the area and a total of 20 habitat features were recorded within The Footprint. Several patches within The Footprint have been mapped as habitat as there were many habitat features in a certain area such as course woody debris and stags. It is recommended that a qualified fauna spotter catcher undertake a pre-clearance survey across the disturbance footprint to identify habitat features prior to clearing and be present during clearing works to check habitat features and relocate fauna, and where possible habitat features in order to minimise impacts to fauna.

ATLAS Stage 3 — EPBC Pre-clearance Report - Hillandale August 2025



Appendix A – Fauna records



| Common Name | Scientific Name |
|---------------------------|-----------------------------|
| Eastern grey kangaroo | Macropus giganteus |
| Red-rumped parrot | Psephotus haematonotus |
| Willie wagtail | Rhipidura leucophrys |
| Masked lapwing | Vanellus miles |
| Apostlebird | Struthidea cinerea |
| Pied butcherbird | Cracticus nigrogularis |
| Galah | Eolophus roseicapilla |
| Australian bustard | Ardeotis australis |
| Crested pigeon | Ocyphaps lophotes |
| Brown falcon | Falco berigora |
| Black-faced cuckoo-shrike | Coracina novaehollandiae |
| Grey-crowned babbler | Pomatostomus temporalis |
| Bar-shouldered dove | Geopelia humeralis |
| Fairy martin | Petrochelidon ariel |
| Double-barred finch | Taeniopygia bichenovii |
| Torresian crow | Corvus orru |
| Welcome swallow | Hirundo neoxena |
| Black-fronted dotterel | Elseyornis melanops |
| Australian grebe | Tachybaptus novaehollandiae |
| Magpie-lark | Grallina cyanoleuca |
| Pied currawong | Strepera graculina |
| Superb fairy-wren | Malurus cyaneus |
| Black-shouldered kite | Elanus axillaris |
| Bearded dragon | Pogona barbata |
| Grey teal | Anas gracilis |
| Australian magpie | Gymnorhina tibicen |
| Wedge-tailed eagle | Aquila audax |
| Red-necked wallaby | Notamacropus rufogriseus |
| Emu | Dromaius novaehollandiae |
| Burns's dragon | Amphibolurus burnsi |
| Sand monitor | Varanus gouldii |

ATLAS Stage 3 — EPBC Pre-clearance Report - Hillandale August 2025



Appendix B – Habitat features



| Habitat type | Habitat location | |
|-------------------------------|-------------------------------------|--|
| Logs | -26.3061136817389, 149.843751325637 | |
| Stag | -26.3104496365203, 149.860117992689 | |
| Hollows | -26.3071343226787, 149.855715838246 | |
| Hollow logs | -26.3032510680815, 149.862784318615 | |
| Logs | -26.3043381625709, 149.863923484141 | |
| Large hollow log | -26.3044931411042, 149.864422852287 | |
| Hollow logs | -26.3065240580668, 149.864886547633 | |
| Scattered logs | -26.3068657066144, 149.86469566733 | |
| Martin mud nests in bank | -26.3074818789267, 149.851112582827 | |
| Pardalote holes/nests | -26.3074885135865, 149.851174244093 | |
| Small stick nest, dome shaped | -26.3106417089938, 149.860083432721 | |
| Scattered logs | -26.3111832600946, 149.860199142088 | |
| Scattered logs | -26.311602862932, 149.860397032926 | |
| Scattered large hollow logs | -26.3105754510471, 149.860658883663 | |
| Small stick nest, dome shaped | -26.3103334655332, 149.860340790162 | |
| Large log stumps | -26.3058305284376, 149.864719857055 | |
| Large log | -26.3108527797986, 149.865412326702 | |
| Coarse woody debris | -26.305664074441, 149.867463815257 | |
| Scattered logs | -26.304678014857, 149.868913368285 | |
| Coarse woody debris | -26.2998971774815, 149.851205101305 | |
| Scattered medium logs | -26.307952518527, 149.864066287931 | |
| Several large logs | -26.3080916594125, 149.859678747656 | |
| Scattered medium logs | -26.3080793318977, 149.864471045498 | |
| Decorticating bark | -26.3084516243952, 149.864287291534 | |
| Scattered medium logs | -26.3084742290864, 149.864376663137 | |
| Scattered medium logs | -26.3086012698115, 149.864387603734 | |
| Medium hollows | -26.3086266864677, 149.864480565005 | |
| Scattered medium logs | -26.3088337471652, 149.864407692786 | |
| Large logs - log pile | -26.3122744793423, 149.881679909253 | |
| Large log | -26.3125006733628, 149.8807684228 | |
| Medium logs | -26.3118587212081, 149.876401409844 | |
| Large log | -26.3125006733628, 149.8807684228 | |
| Medium logs | -26.3118587212081, 149.876401409844 | |
| Coarse woody debris | -26.3080875505359, 149.864032239098 | |
| Large log | -26.3080560281676, 149.864224086021 | |
| Hollow log | -26.3080322155115, 149.863858566882 | |
| Log | -26.3081463314173, 149.862736960856 | |
| | | |

ATLAS Stage 3 — EPBC Pre-clearance Report - Hillandale August 2025



| Habitat type | Habitat location |
|-----------------------------------|-------------------------------------|
| Two stags with peeling bark | -26.3082116950978, 149.864375263533 |
| Hollow | -26.3091104547951, 149.864601291458 |
| Two hollow stags | -26.3121378478342, 149.876345390029 |
| Large log in gully | -26.3112199322561, 149.875303672875 |
| Small stick nest | -26.3111721769088, 149.875425680268 |
| Medium logs in gully | -26.3113646339101, 149.875260923859 |
| Scattered medium and large logs | -26.3102743220926, 149.867308670402 |
| Scattered medium and large logs | -26.3104818332987, 149.866163735136 |
| Scattered medium and large logs | -26.3104773520873, 149.865808353487 |
| Logs | -26.3264139793092, 149.858409266513 |
| Logs | -26.3262445100708, 149.858366941468 |
| Burrow - Varanus gouldii observed | -26.3266500182459, 149.85897171078 |
| Scattered CWD | -26.3255992851252, 149.857856073036 |
| Scattered large logs | -26.3247622952369, 149.857884964565 |
| Small stag with cracks | -26.324532270956, 149.857841075545 |
| Two medium logs | -26.3269915759571, 149.859753439232 |

ATLAS Stage 3 — EPBC Pre-clearance Report - Hillandale August 2025



Appendix C – Weed species list



| Scientific Name | Common Name | Status | Abundance |
|------------------------|-----------------------|------------------------|------------|
| Bidens pilosa | Cobbler's pegs | * | Rare |
| Chloris gayana | Rhodes grass | * | Occasional |
| Cirsium vulgare | Spear thistle | * | Rare |
| Eragrostis trichophora | Hairyflower lovegrass | * | Frequent |
| Erigeron bonariensis | Flaxleaf fleabane | * | Common |
| Glandularia aristigera | Desert verbena | * | Occasional |
| Gomphrena celosioides | Gomphrena weed | * | Frequent |
| Megathyrsus maximus | Guinea grass | * | Occasional |
| Melinis repens | Natal grass | * | Occasional |
| Opuntia stricta | Common prickly pear | Restricted-Category- | Occasional |
| Opuntia tomentosa | Velvety tree pear | Restricted- Category 3 | Frequent |
| Senna occidentalis | Coffee senna | * | Rare |
| Setaria pumila | Pale pigeon grass | * | Occasional |
| Sida rhombifolia | Common sida | * | Occasional |
| Vachellia farnesiana | Mimosa bush | * | Rare |
| Verbesina encelioides | Crownbeard | * | Rare |
| Xanthium occidentale | Noogoora burr | * | Rare |
| Zinnia peruviana | Wild zinnia | * | Rare |

ATLAS Stage 3 — EPBC Pre-clearance Report - Hillandale August 2025



Appendix D – Erosion points and photos



| Type of erosion | GPS | Photo of erosion |
|----------------------------|------------------------|------------------|
| Gully erosion | -26.312197, 149.867154 | |
| Beginning of erosion gully | -26.309866, 149.851060 | N/A |
| Gully erosion | -26.321265, 149.861284 | |
| Gully erosion | -26.317769, 149.861049 | |



| Type of erosion | GPS | Photo of erosion |
|----------------------------|------------------------|------------------|
| Gully erosion | -26.311777, 149.878461 | |
| Gully erosion | -26.311955, 149.876855 | |
| Sheet erosion | -26.319381, 149.855936 | |
| Beginning of erosion gully | -26.305729, 149.842739 | N/A |